

## Use of Real-Time PCR Technique in Determination of Major Fibrolytic and non Fibrolytic Bacteria Present in Indian Surti Buffaloes (*Bubalus bubalis*)

KRISHNA M. SINGH<sup>1,\*</sup>, AJAI K. TRIPATHI<sup>1</sup>, PARESH R. PANDYA<sup>2</sup>, SUBHASH PARNERKAR<sup>2</sup>,  
DHARAMSI N. RANK<sup>3</sup>, RAMESH K. KOTHARI<sup>4</sup> and CHAITANYA G. JOSHI<sup>1</sup>

<sup>1</sup>Department of Animal Biotechnology, College of Veterinary Science and Animal Husbandry (CVSAH),  
Anand Agricultural University (AAU), Anand (388 001), Gujarat, India

<sup>2</sup>Animal Nutrition Research Station, CVSAH, AAU, Anand, Gujarat, India

<sup>3</sup>Department of Animal Genetics & Breeding, CVSAH, AAU, Anand, Gujarat, India

<sup>4</sup>Department of Microbiology, Christ College, Rajkot, Gujarat, India

<sup>5</sup>Department of Genetics, ARIBAS, New V V Nagar, Anand, India (Present address)

Received .... 2013, revised ... 2013, accepted .... 2013

### Abstract

In the milk industry in India, buffalo breeds are most commonly used for milk production. Efficiency of fiber digestion in ruminants is critical for animal productivity. Bacteria play an important role in fiber digestion and utilization. Absolute quantification real-time PCR was used to quantify ten bacterial species in rumen fluid of Surti buffalo fed green fodder, dry roughage and compound concentrate mixture. Abundance of each target taxon was calculated as a fraction of the total 16S rRNA gene copies in the samples, using taxon-specific primers. Bacterial populations showed a clear predominance of *Ruminococcus albus*, which comprised 5.66% of the bacterial rRNA gene copies in the samples. However, only 0.9% to 4.24% of the bacterial rRNA gene copies were represented by the ruminal *Fibrobacter succinogenes*, *Ruminococcus flavefaciens* and *Prevotella* species. The proportion of rRNA gene copies attributable to *Selenomonas ruminantium*, *Streptococcus bovis*, *Ruminobacter amylophilus*, *Treponema bryantii* and *Anaerovibrio lipolytica* was even less abundant, each comprising <0.11% of the bacterial rRNA gene copies. The data suggest that the aggregate abundance of the most intensively studied ruminal bacterial species is relatively low and that a large fraction of the uncultured population represents a single bacterial genus.

**Key words:** Surti buffalo; fibrolytic; non-fibrolytic bacteria; real time PCR

### Introduction

The digestion of plant material and subsequent conversion for energy requirements to the host ruminant are performed through a complex symbiotic relationship of microbiota within the rumen (Mackie 1997). The composition and proportion of microorganisms are influenced by external factors, such as diet, feeding frequency, age, geographical location and ruminant-host interaction (Hungate 1966). Bacteria are considered to be the most important for the biological degradation of dietary fibers due to their fibrolytic activity and biomass in the rumen. Although fibrolytic species such as *Fibrobacter succinogenes*, *Ruminococcus albus* and *Ruminococcus flavefaciens* play a key role in plant fiber degradation (Forsberg *et al.*, 1997), positive interactions between them and non-fibrolytic bacteria such as *Selenomonas ruminantium* and *Treponema bryantii* have been reported (Kudo *et al.*, 1987). In an early study, the synergism between *R. flavefaciens* and

*S. ruminantium* was suggested as enhancing propionate production (Scheifinger and Wolin 1973). Sawanon and Kobayashi, (2006) reported that fiber digestibility and propionate production significantly increased in coculture of *R. flavefaciens* and *S. ruminantium* compared to mono-culture of *R. flavefaciens*. These findings indicate that non-fibrolytic bacteria may also be important to facilitate plant fiber degradation in the rumen. Therefore, non-fibrolytic bacteria as well as fibrolytic bacteria should be monitored in order to estimate overall contribution of bacteria to ruminal fiber digestion.

India possesses more than 50% of world's buffalo population; Indian buffaloes produce more than 60% milk in India (Kumar *et al.*, 2007). Surti is a popular breed of buffalo found in central Gujarat state. The Surti buffaloes are of medium size and docile temperament and body weight 350 to 375 kg at maturity.

Increased knowledge concerning the rumen fibrolytic and non fibrolytic bacterial population will allow insight into the fiber-digestion capabilities of ruminant

\* Correspondence: Singh K.M. phone/fax: +91 2692 229189; e-mail: Kmsingh18@gmail.com

animals. Our animals mainly sustain on crop residues, the bacterial population is expected much different than that of HF cattle. However, very limited research has been conducted in Indian buffalo with regard to the ruminal microbial population using molecular techniques. Therefore, this study was conducted to determine the population size of rumen bacteria in Surti buffalo by real-time PCR assays fed green fodder Napier bajra, pasture grass (*Dichanthium annulatum*), and compound concentrate mixture.

## Experimental

### Materials and Methods

**Animals, diet and collection of rumen fluid.** The experiment was carried out on three adult Surti buffaloes reared at the Department of Animal Nutrition, College of Veterinary Science and A.H., Anand. All the animals were maintained under uniform feeding regime for minimum 30 days. The diet was consisting of green fodder Napier bajra 21 (*Pennisetum purpureum*), mature pasture grass (*Dichanthium annulatum*) and

compound concentrate mixture (20% crude protein, 65% total digestible nutrients). Animals were offered 10 kg green fodder, 6 kg dry fodder and 2.0 kg of concentrate mixture once a day. All Animals were let loose daily for 2 hrs morning and evening, during which they had free access to drinking water. Samples of rumen liquor (about 500 ml) were collected at 4 hrs, 8 hrs and 24 hrs after feeding by a suction pump using a flexible stomach tube (Khamapa *et al.*, 2006).

**DNA extraction and PCR amplification of 16S RNA genes.** Total DNA was extracted separately by using a commercially available kit according to the manufacturer's instructions (QIAGEN Stool kit; QIAGEN, CA). The total DNA mixture was used as a template in PCR to amplify 16S rDNA. The target DNA of total bacteria, bovine bacteroides, fibrolytic and non fibrolytic bacteria were amplified from the metagenomic DNA, as described previously by (Muyzer *et al.*, 1993; Layton *et al.*, 2006; Tajima *et al.*, 2001; Koike and Kobayashi 2001).

**Preparation of standard plasmid for real-time PCR assays.** After the confirmation of a single band of the correct size with respective pair of primers (Table I)

Table I  
Primers for real time PCR assay and PCR conditions

Target	Primer Sequence (5'-3')	Annealing temp. (°C)	Product size (bp)	Reference
<b>Total bacteria</b>	P1-CCTACGGGAGGCAGCAG P2-ATTACCGCGGCTGCTGG	60	194	(Muyzer <i>et al.</i> , 1993)
<b>Bacteroides (bovine)</b>	P1-GAAG(G/A)CTGAACCAGCCAAGTA P2-GCTTATTCATACGGTACATACAAG	57	100	(Layton <i>et al.</i> , 2006)
<b>Fibrolytic bacteria</b>				
<i>Fibrobacter succinogenes</i>	P1-GGTATGGGATGAGCTTGC P2-GCCTGCCCCTGAACATATC	60	446	(Tajima <i>et al.</i> , 2001)
<i>Ruminococcus albus</i>	P1-CCCTAAAAGCAGTCTTAGTTTCG P2-CCTCCTTGCGGTTAGAACA	60	175	(Koike and Kobayashi, 2001)
<i>Ruminococcus flavefaciens</i>	P1-GGACGATAATGACGGTACTT P2-GCAATC(CT)GAACTGGGACAAT	55	295	(Koike and Kobayashi, 2001)
<i>Prevotella bryantii</i>	P1-AGTCGAGCGGTAAGATTG P2-CAAAGCGTTTCTCTCACT	68	540	(Tajima <i>et al.</i> , 2001)
<i>Prevotella ruminicola</i>	P1-GGTTATCTTGAGTGAGTT P2-CTGATGGCAACTAAAGAA	53	485	(Tajima <i>et al.</i> , 2001)
<b>Non-fibrolytic bacteria</b>				
<i>Anaerovibrio lipolytica</i>	P1-TGGGTGTTAGAAATGGATTTC P2-CTCTCCTGCACTCAAGAATT	57	597	(Tajima <i>et al.</i> , 2001)
<i>Ruminobacter amylophilus</i>	P1-CAACCAGTCGCATTTCAGA P2-CACTACTCATGGCAACAT	57	642	(Tajima <i>et al.</i> , 2001)
<i>Selenomonas ruminantium</i>	P1-TGCTAATACCGAATGTTG P2-TCCTGCACTCAAGAAAGA	57	513	(Tajima <i>et al.</i> , 2001)
<i>Streptococcus bovis</i>	P1-CTAATACCGCATAACAGCAT P2-AGAAACTTCCTATCTCTAGG	57	869	(Tajima <i>et al.</i> , 2001)
<i>Treponema bryantii</i>	P1-ACTGCAGCGCGAACTGTCAGA P2-ACCTTACGGTGGCAGTGTCTC	57	412	(Tajima <i>et al.</i> , 2001)

on an agarose gel, the PCR products were excised from the gel. The PCR products were purified using the Qiagen gel Purification Kit (Qiagen, CA), and then ligated into pTZR57T/A cloning vector (Fermentas, UK). The ligated products were transformed to competent *Escherichia coli* DH5 alpha cells by heat shock. Plasmids were purified from positive clones using a QIAprep spin miniprep kit (Qiagen, Valencia, CA, USA), and the plasmids containing the correct insert were screened out by PCR amplification with respective primer sets. The concentration of the plasmid was determined with a Nanodrop spectrophotometer. Copy number of each standard plasmid was calculated using formula; Copy No/ $\mu$ l = Concentration of plasmids (gm/ $\mu$ l)  $\times$   $6.022 \times 10^{23}$  / length of recombinant plasmid (bp)  $\times$  660, (660 = Molecular weight of nucleotide base,  $6.022 \times 10^{23}$  = Avogadro's number). Ten-fold dilution series ranging from  $10^9$  to 10 copies were prepared for each target. Real-time PCR was performed with ABI system (ABI 7500). The Qiagen DNA Master SYBR Green I was used for PCR reaction. The optimal amplification conditions for each primer set were obtained with 10 mol each primer with the combination of annealing temperature shown in Table I. The 10-fold dilution series of the standard plasmid for the respective target was run along with the corresponding samples in duplicate. The copy numbers of 16S rRNA genes of targeted microorganisms were calculated using the following equation:  $(QM \times C \times DV) / (S \times V)$ , where QM was the quantitative mean of the copy number C was the DNA concentration of each sample, DV was dilution volume of extracted DNA, S was the DNA amount (ng) subjected to analysis and V was rumen fluid volume subjected to DNA extraction. In the reaction for all standard, nearly perfect linear regressions ( $r^2 = 0.9930$  to  $0.9995$ ), intercept (31.26 to 44.78) and slope ( $-3.2$  to  $-4$ ) were obtained between threshold cycle and quantities of standard for all targets and data generated from the reaction were used for further analysis.

## Results and Discussion

In the present study, we assessed the rumen bacterial species, using real-time PCR assays. The results of qualitative PCR detection of fibrolytic bacteria and non fibrolytic bacteria are given in figure 1. Real-time PCR-based quantification of representative rumen bacteria have been also reported in other rumen ecology studies (Ozutsumi *et al.*, 2006).

**Quantitation of total bacteria, fibrolytic and non fibrolytic bacteria.** Table I shows the population sizes of total bacteria and the target species in Surti buffalo rumen. Total bacteria and bovine bacteroides were detected at  $5.8 \times 10^9$  copies and  $3.1 \times 10^8$  per ml ruminal




Fig. 1. Qualitative PCR detection of fibrolytic bacteria and non-fibrolytic bacteria. Lane: 1, *Ruminococcus albus*; 2, *Fibrobacter succinogenes*; 3, *Ruminococcus flavefaciens*; 4, *Ruminobacter amylophilus*; 5, *Streptococcus bovis*; 6, *Prevotella bryantii*; 7, *Anaerovibrio lipolytica*; 8, *Prevotella ruminicola*; 9, *Selenomonas ruminantium*; 10, *Treponema bryantii*; 11, Total bacteria; 12, bacteroides; Lane M, DNA size marker.

fluid after 4 hrs incubation in the rumen and increased to  $5.9 \times 10^9$  and  $3.2 \times 10^8$  copies per ml ruminal fluid at 24 hrs respectively. Among the fibrolytic bacteria, *R. albus* was most abundantly detected ( $3.0 \times 10^8$  copies/ ml of rumen fluid accounting 5.66% of 16S r RNA gene copies after 24 hrs and followed by *R. flavefaciens* ( $2.5 \times 10^8$  copies/ml of rumen fluid, corresponding to 4.24% of 16S r RNA gene copies (Table II). In non fibrolytic bacteria, *Streptococcus bovis* ( $5.3 \times 10^6$  copies/ml of rumen fluid) and *Selenomonas ruminantium* ( $1.1 \times 10^6$  copies/ml of rumen fluid) was detected, accounting for 0.11% and 0.025% of 16S r RNA gene copies at 24 hrs (Table II). The population size of all fibrolytic and nonfibrolytic bacterial species were significantly higher 24 hrs after feeding except *S. ruminantium*, *R. amylophilus* and *T. bryantii*.

To our knowledge, no previous study has reported the population size of rumen microbes of Indian buffalo, particular in the Surti buffalo rumen. In the present study, our observation indicates that a large number of bacteria were present in the ruminal fluid. Because each bacterial species has a different copy number of 16S rRNA gene (ranging from 1 to 15; (Klappenbach *et al.*, 2001), this is comparable to the level of rumen bacteria counted by microscopy in the literature (Hungate, 1966).

The present results among fibrolytic bacteria revealed that *R. albus*, *R. flavefaciens* and *F. succinogenes* are a core member of bacteria, playing a significant and important role in the digestion of fiber rich diet. *R. albus* was most dominant ( $3.0 \times 10^8$  copies/ml of rumen fluid) among the three species, followed by *R. flavefaciens* ( $2.5 \times 10^8$  copies/ml of rumen fluid) and *F. succinogenes* ( $1 \times 10^8$  copies/ml of rumen fluid). Similarly, Ozutsumi *et al.*, (2006) reported that the numbers of *R. albus* ( $\log_{10} 8.3 \pm 6.3$  per ml) and *R. flavefaciens* ( $\log_{10} 8.8 \pm 7.9$  per ml) were higher than *F. succinogenes* ( $\log_{10}$

Table II  
Distribution of microbial population size and proportions of total bacteria to ruminal fluid of Surti buffalo

Targets	Copy of target DNA/ml of ruminal fluid			% Total 16S r RNA gene		
	4 hrs	8 hrs	24 hrs	4 hrs	8 hrs	24 hrs
Total bacteria	$5.8 \times 10^9$	$4.5 \times 10^9$	$5.9 \times 10^9$	–	–	–
Bovine bacteroides	$3.1 \times 10^8$	$2.3 \times 10^8$	$3.2 \times 10^8$	–	–	–
<i>Ruminococcus albus</i>	$1.37 \times 10^8$	$7.3 \times 10^7$	$3.0 \times 10^8$	2.36	1.62	5.66
<i>Fibrobacter succinogenes</i>	$7.2 \times 10^7$	$2.8 \times 10^6$	$2.5 \times 10^8$	1.24	0.06	4.24
<i>Ruminococcus flavefaciens</i>	$3.5 \times 10^7$	$2.8 \times 10^7$	$1.0 \times 10^8$	0.60	0.62	1.72
<i>Prevotella bryantii</i>	$2.5 \times 10^3$	$1.6 \times 10^3$	$1.0 \times 10^4$	0.000043	0.000036	0.000017
<i>Prevotella ruminicola</i>	$3.6 \times 10^7$	$4.2 \times 10^7$	$5.7 \times 10^7$	0.62	0.93	0.97
<b>Non fibrolytic bacteria</b>						
<i>Streptococcus bovis</i>	$1.2 \times 10^4$	$7.4 \times 10^3$	$6.3 \times 10^6$	0.00021	0.00016	0.11
<i>Selenomonas ruminantium</i>	$9.5 \times 10^5$	$9.5 \times 10^5$	$1.5 \times 10^6$	0.016	0.021	0.025
<i>Ruminobacter amylophilus</i>	$1.6 \times 10^6$	$2.2 \times 10^6$	$1.6 \times 10^6$	0.028	0.049	0.0034
<i>Treponema bryantii</i>	$1.9 \times 10^5$	$3.2 \times 10^5$	$1.3 \times 10^5$	0.0033	0.0071	0.0022
<i>Anaerovibrio lipolytica</i>	$2.7 \times 10^3$	$8.0 \times 10^3$	$9.9 \times 10^3$	0.000047	0.00018	0.00017

7.7 ± 6.8 per ml) in the unfauinated cattle rumen fed 66% Sudan grass hay and 34% concentrate mixture. However, Wanapat and Cherdthong (2009) reported that *F. succinogenes* was the major cellulolytic bacterium of rumen digesta in the swamp buffalo and was present  $2.65 \times 10^9$  to  $3.54 \times 10^9$  copies/ml in ruminal fluid. Koike and Kobayashi (2001) also reported that *F. succinogenes* was the major cellulolytic bacteria of rumen digesta in sheep and present at only 0.1% of total population and that ruminococci were relatively minor. The scarcities of the two ruminococci were surprising, considering that they were representative cellulolytics (*i.e.* ruminal densities ranging from 0.1% (Stahl *et al.*, 1998) to 6.6% (Briesacher *et al.*, 1992) for *F. succinogenes*, and from 1.3% to 2.9% for *Ruminococcus* ssp. (Krause *et al.*, 1999). Ruminal *prevotella* are known to possess oligo-saccharolytic and xylanolytic activities and to occupy the ecological niches of the second line degraders. The density of *P. ruminicola* was 0.97% of total bacteria at 24 hrs (Table II). The quantity of *P. bryantii* was little low demonstrating opposite kinetics, suggesting its role in starch degradation. Dominance of *Prevotella* and low abundance of classical ruminal bacterial species have been also reported by Stevenson and Weimer, (2009) in the bovine rumen.

Non-fibrolytic bacteria such as *S. ruminantium* and *T. bryantii* were detected in the fiber-associated community, using comparative 16S rRNA gene analysis (Koike *et al.*, 2003). The present study quantitatively confirmed the nonfibrolytic bacteria to fed green fodder *Pennisetum purpureum* and *Dichanthium annulatum* (Table II). In particular, *S. bovis*, *S. ruminantium* and *Ruminobacter amylophilus* had the highest proportion among the non fibrolytic bacterial species assessed in the present study. This finding suggests the

fiber-attachment ability of *S. ruminantium*. Minato and Suto (1978) indicated that some of the non-fibrolytic bacteria possessed the ability to attach to cellulose at a similar extent to that of fibrolytic bacteria. In addition, non-fibrolytic species such as *Selenomonas* ssp. and *Streptococcus* ssp. were isolated from plant cell material in rumen digesta (Cheng *et al.*, 1984). Attachment of non-fibrolytic bacteria to green fodder may be mediated by glycocalyxes (Minato *et al.*, 1993), which are commonly found in rumen bacteria during their colonization in plant cell wall (Cheng *et al.*, 1980). These reports support the non-fibrolytic bacteria in ruminal fluid of Surti buffalo as observed in the present study. In such a relationship, fibrolytic bacteria provide the hydrolyzed product from cellulose to non-fibrolytic bacteria, while non-fibrolytic bacteria indirectly facilitate fiber degradation by preventing the accumulation of bacterial metabolites such as succinate and cellodextrins (Wolin *et al.*, 1997). Utilization of cellodextrins from cellulose hydrolysis is crucial in further fiber digestion (Russell 1985), because celluloses are highly sensitive to feedback inhibition by cellobiose, which is confirmed in *F. succinogenes* (Maglione *et al.*, 1997). In the present study, the population sizes of *S. bovis* and *S. ruminantium* in fluid were greater than those of the other non-fibrolytics (Table II). After 24 hrs ruminal incubation, they were estimated to be 0.11 and 0.025% of total bacteria, respectively (Table II). This result indicates that these two species were more numbers and it is known that *S. bovis* and *S. ruminantium* are able to utilize hydrolysis product of polysaccharide such as cellodextrins and maltodextrins for their growth (Cotta, 1992). Therefore, *S. bovis* and *S. ruminantium* in rumen might grow by utilizing the maltodextrins released during starch degradation (Nouaille *et al.*, 2005).



*Treponema* spp. are a commonly detected bacterial group in the rumen that are involved in the degradation of soluble fibers. Bekele *et al.*, (2011) also reported high level of treponema group in sheep rumen samples. Moreover, this fact strongly highlights the importance of non-fibrolytic bacteria for fiber digestion. Culture-independent molecular studies clearly indicate an abundance of uncultured bacteria, protozoa and archaea in the rumen (Shin *et al.*, 2004). The sum of proportions of 10 representative bacterial species was calculated to be 12.73% of total bacteria. This result strongly suggests the significance of uncultured bacteria in plant fiber degradation. Therefore, it is necessary to perform quantitative determination of the community structure of rumen microbes including the uncultured microbes and also to cultivate such microbe.

In conclusion, major representative groups of ruminal microbes covering fibrolytic and non-fibrolytic bacterial communities were detected. Results revealed that *R. albus* was the most dominant fibrolytic among the all detected fibrolytic species. Non-fibrolytic *S. bovis* and *S. ruminantium* were also detected with abundance in rumen fluid. The high magnitude of non-fibrolytic group on the plant fiber suggests the development of mutual relationships between fibrolytic and non-fibrolytic bacterial communities. Present study indicates that rumen system of Surti buffalo is harbours the diverse group of microbial communities with high population densities.

#### Acknowledgement

Financial support provided by the Department of Biotechnology, Govt. of India, New Delhi to conduct the study reported here is acknowledged with respect and gratitude.

#### Literature

- Bekele A.Z., S. Koike and Y. Kobayashi. 2011. Phylogenetic diversity and dietary association of rumen *Treponema* revealed using group-specific 16S rRNA gene-based analysis. *FEMS Microbiol. Lett.* 316(1): 51–60.
- Briesacher S.L., T. May, K.N. Grigsby, M.S. Kerley, R.V. Anthony and J.A. Paterson. 1992. Use of DNA probes to monitor nutritional effects on ruminal prokaryotes and *Fibrobacter succinogenes* S85. *J. Anim. Sci.* 70: 289–295.
- Cheng K.J., C.S. Stewart, D. Dinsdale and J. Costerton. 1984. Electron microscopy of bacteria involved in the digestion of plant cell wall. *An. Feed. Sci. Tech.* 10: 93–120.
- Cheng K., J.P. Fay, R.E. Howarth and J.W. Costerton. 1980. Sequence events in the digestion of fresh legume leaves by rumen bacteria. *Appl. Environ. Microbiol.* 40: 613–625.
- Cotta M.A. 1992. Interaction of ruminal bacteria in the production and utilization of maltooligosaccharides from starch. *Appl. Environ. Microbiol.* 58: 48–54.
- Denman S.E. and C.S. McSweeney. 2006. Development of a real-time PCR assay for monitoring anaerobic fungal and cellulolytic bacterial populations within the rumen. *FEMS Microbiology Ecology* 85(3): 572–582.
- Forsberg C.W., K. Cheng and B.A. White. 1997. Polysaccharide degradation in the rumen and large intestine. In: Mackie R.I., White B.A. (eds), *Gastrointestinal Microbiology*. Chapman & Hall, New York, pp. 319–379.
- Hungate R.E. 1966. The Rumen and its Microbes. *Academic Press*, New York.
- Khamapa S., M. Wanapat, C. Wachirapakorn and N. Nontaso. 2006. Effects of urea level and sodium di-malate in concentrate containing high cassava chip on ruminal fermentation efficiency, microbial protein synthesis in lactating dairy cows raised under tropical condition. *Asian- Aust. J. Anim. Sci.* 19: 837–844.
- Klappenbach J.A., P.R. Saxman, J.R. Cole and T.M. Schmidt. 2001. rrndb: The Ribosomal RNA Operon Copy Number Database. *Nucl. Acids. Res.* 29: 181–184.
- Koike S. and Y. Kobayashi. 2001. Development and use of competitive PCR assays for the rumen cellulolytic bacteria: *Fibrobacter succinogenes*, *Ruminococcus albus* and *Ruminococcus flavefaciens*. *FEMS. Microbiol. Lett.* 204: 361–366.
- Koike S., J. Pan, Y. Kobayashi and K. Tanaka. 2003. Kinetics of in sacco fiber-attachment of representative ruminal cellulolytic bacteria monitored by competitive PCR. *J. Dairy Sci.* 86: 1429–1435.
- Koike S., H. Yabuki and Y. Kobayashi. 2007. Validation and application of real-time polymerase chain reaction assays for representative rumen bacteria. *Anim. Sci. J.* 78: 135–141.
- Krause D.O., B.P. Dalrymple, W.J. Smith, R.I. Mackie and C.S. McSweeney. 1999. 16S rDNA sequencing of *Ruminococcus albus* and *Ruminococcus flavefaciens*: design of a signature probe and its application in adult sheep. *Microbiol.* 145: 1797–1807.
- Kudo H., K.J. Cheng and J.W. Costerton. 1987. Interactions between *Treponema bryantii* and cellulolytic bacteria in the in vitro degradation of straw cellulose. *Can. J. Microbiol.* 33: 244–248.
- Kumar S., M. Nagarajan, J. Sandhu, N. Kumar and V. Behl. 2007. Phylogeography and domestication of Indian river buffalo. *BMC. Evol. Biol.* 7, 186.
- Layton A., L. McKay, D. Williams, V. Garrett, R. Gentry and G. Sayler. 2006. Development of Bacteroides 16S rRNA Gene Taq-Man-Based Real-Time PCR Assays for Estimation of Total, Human, and Bovine Fecal Pollution in Water. *Appl. Environ. Microbiol.* 72 (6): 4214–4224.
- Leser T.D., M. Boye and N.B. Hendriksen. 1995. Survival and activity of a *Pseudomonas* sp. strain B13 (FR1) in a marine microcosm determined by quantitative PCR and an rRNA-targeting probe and its effect on the indigenous bacterioplankton. *Appl. Environ. Microbiol.* 61: 1201–1207.
- Lin C., B. Flesher, W.C. Capman, R.I. Amann and D.A. Stahl. 1994. Taxon specific hybridization probes for fibre-digesting bacteria suggest novel gut-associated *Fibrobacter*. *Syst. Appl. Microbiol.* 17: 418–424.
- Mackie R.I. 1997. Gut environment and evolution of mutualistic fermentative digestion. *Gastrointestinal Microbiology*, Vol. 1 (Mackie R.I. & White B.A., eds), Chapman & Hall, New York, pp13–35.
- Maglione G., J.B. Russell and D.B. Wilson. 1997. Kinetics of cellulose digestion by *Fibrobacter succinogenes* S85. *Appl. Environ. Microbiol.* 63: 665–669.
- Minato H., M. Mitsumori and K.J. Cheng. 1993. Attachment of microorganisms to solid substrate in the rumen. In: Shimada K, Hoshino S (eds), *Genetics, Biochemistry and Ecology of Lignocellulose Degradation*. *Uni Publishers. Tokyo*. pp, 139–145.
- Minato H. and T. Suto. 1978. Technique for fractionation of bacteria in rumen microbial ecosystem. II. Attachment of bacteria isolated from bovine rumen to cellulose powder in vitro and elution of bacteria attached there from. *J. Gen. Appl. Microbiol.* 24: 1–16.
- Mosoni P., F. Chaucheyras-Durand, C. Béra-Maillet and E. Forano. 2007. Quantification by real-time PCR of cellulolytic bacteria in the rumen of sheep after supplementation of a forage diet with readily

- fermentable carbohydrates: effect of a yeast additive. *J. Appl. Microbiol.* 103(6): 2676–2685.
- Muyzer G., E.C. Dewall and A.G. Uitterlinden.** 1993. Profiling of complex microbial populations by denaturing gradient gel electrophoresis analysis of polymerase chain reaction amplified genes coding for 16S rRNA. *Appl. Environ. Microbiol.* 59: 695–700.
- Nouaille R., M. Matulova, A.M. Delort and F. Forano.** 2005. Oligosaccharide synthesis in *Fibrobacter succinogenes* S85 and its modulation by the substrate. *FEBS Journal.* 272: 2416–2427.
- Ozutsumi Y., K. Tajima, A. Takenaka and H. Itabashi.** 2006. Real time PCR detection of the effect of protozoa on rumen bacteria in cattle. *Curr. Microbiol.* 52: 158–162.
- Reilly K. and Attwood GT.** 1998. Detection of *Clostridium proteoclasticum* and closely related strains in the rumen by competitive PCR. *Appl. Environ. Microbiol.* 64: 907–913.
- Russell J.B.** 1985. Fermentation of cellodextrins by cellulolytic and noncellulolytic rumen bacteria. *Appl. Environ. Microbiol.* 49: 572–576.
- Sawanon S. and Y. Kobayashi.** 2006. Synergistic fibrolysis in the rumen by cellulolytic *Ruminococcus flavefaciens* and noncellulolytic *Selenomonas ruminantium*: evidence in defined cultures. *Anim. Sci. J.* 77: 208–214.
- Scheifinger C.C. and M.J. Wolin.** 1973. Propionate formation from cellulose and soluble sugars by combined cultures of *Bacteroides succinogenes* and *Selenomonas ruminantium*. *J. Appl. Microbiol.* 26: 89–795.
- Shinkai T., T. Ueki and Y. Kobayashi.** 2010. Detection and identification of rumen bacteria constituting a fibrolytic consortium dominated by *Fibrobacter succinogenes*. *Animal Science Journal.* 81: 72–79.
- Shin E.C., K.M. Cho, W.J. Lim, S.Y. Hong, C.L. An, E.J. Kim, Y.K. Kim, B.R. Choi, J.M. An, J.M. Kang, H. Kim and H.D. Yun.** 2004. Phylogenetic analysis of protozoa in the rumen contents of cow based on the 18S rDNA sequences. *J. Appl. Microbiol.* 97: 378–383.
- Stahl D.A., B. Flesher, H.R. Mansfield and L. Montgomery.** 1988. Use of phylogenetically based hybridization probes for studies of ruminal microbial ecology. *Appl. Environ. Microbiol.* 54: 1079–1084.
- Stevenson D.M. and P.J. Weimer.** 2007. Dominance of Prevotella and low abundance of classical ruminal bacterial species in the bovine rumen revealed by relative quantification real-time PCR. *Appl. Microbiol. Biotechnol.* 75(1): 165–74
- Stiverson J., M. Morrison and Z. Yu.** 2011. Populations of Select Cultured and Uncultured Bacteria in the Rumen of Sheep and the Effect of Diets and Ruminant Fractions. *International Journal of Microbiology* doi:10.1155/2011/750613.
- Sylvester J.T., S.R. Karnati, Z. Yu, M. Morrison and J.L. Firkins.** 2004. Development of an assay to quantify rumen ciliate protozoal biomass in cows using real-time PCR. *J. Nutr.* 134: 3378–3384.
- Tajima K., R.I. Aminov, T. Nagamine, H. Matsui, M. Nakamura and Y. Benno.** 2001. Diet-dependent shifts in the bacterial population of the rumen revealed with real-time PCR. *Appl. Environ. Microbiol.* 67: 2766–2774.
- Wanapat M. and A. Cherdthong.** 2009. Use of Real-Time PCR Technique in Studying Rumen Cellulolytic Bacteria Population as Affected by Level of Roughage in Swamp Buffalo. *Curr. Microbiol.* 58: 294–299.
- Wolin M.J., T.L. Miller and C.S. Stewart.** 1997. Microbe-microbe interactions. In: Hobson PN, Stewart CS (eds). *The Rumen Microbial Ecosystem*, Blackie Academic and Professional London. Pp. 467–491.
- Zimmermann K. and J.W. Mannhalter.** 1996. Technical aspects of quantitative competitive PCR. *Bio.Tech.* 21: 268–279.